



Journal of Cardiology

journal homepage: www.elsevier.com/locate/jjcc

Letter to the Editor

Pivotal role of elevated heart rate in the cardiovascular continuum

We congratulate Custodis et al. for demonstrating that elevated heart rate (HR) plays a pivotal role in the vascular pathophysiology of the cardiovascular continuum [1]. An elevated HR promotes atherosclerosis via not only sympathetic overactivation, but also by modulating the local hemodynamic environment. An elevated HR increases the exposure of the endothelium to systolic low shear stress (SS), attenuates the diastolic SS, and increases local tensile stress, which change the arterial structure in patients with cardiovascular risks. Thereby, HR lowering might act specifically on atherosclerosis-prone lesions, contributing to a residual risk reduction [2]. The prognostic benefit of β blockers in patients with stable coronary artery disease has been called into question [3]. These findings might be due in part to inadequate HR control [4], indicating that the physician did not use a β blocker in a HR-guided manner, because no optimal HR level is described in the guideline documentation. The SIGNIFY study will reveal the prognostic significance of HR lowering for stable CAD patients [5].

References

- [1] Custodis F, Reil JC, Laufs U, Bohm M. Heart rate: a global target for cardiovascular disease and therapy along the cardiovascular disease continuum. *J Cardiol* 2013;62:183–7.
- [2] Inoue T, Iseki K, Ohya Y. Heart rate as a possible therapeutic guide for the prevention of cardiovascular disease. *Hypertens Res* 2013;36:838–44.
- [3] Bangalore S, Steg G, Deedwania P, Crowley K, Eagle KA, Goto S, Ohman EM, Cannon CP, Smith SC, Zeymer U, Hoffman EB, Messerli FH, Bhatt DL, REACH Registry Investigators. β -Blocker use and clinical outcomes in stable outpatients with and without coronary artery disease. *JAMA* 2012;308:1340–9.
- [4] Steg PG, Ferrari R, Ford I, Greenlaw N, Tardif JC, Tendera M, Abergel H, Fox KM, CLARIFY Investigators. Heart rate and use of beta-blockers in stable outpatients with coronary artery disease. *PLoS ONE* 2012;7:e36284.
- [5] Fox K, Ford I, Steg PG, Tardif JC, Tendera M, Ferrari R. Rationale, design, and baseline characteristics of the study assessing the morbidity–mortality benefits of the if inhibitor ivabradine in patients with coronary artery disease (SIGNIFY trial): a randomized, double-blind, placebo-controlled trial of ivabradine in patients with stable coronary artery disease without clinical heart failure. *Am Heart J* 2013;166, 654–661.e6.

Taku Inoue (MD, PhD)*

Center of Residency and Fellowship Program,
University Hospital of the Ryukyus, 207 Uehara,
Nishihara, Nakagami-gun, Okinawa 903-0125, Japan

Kunitoshi Iseki (MD, PhD)

Dialysis Unit, University Hospital of the Ryukyus,
207 Uehara, Nishihara, Nakagami-gun,
Okinawa 903-0125, Japan

* Corresponding author. Tel.: +81 98 895 1195.
E-mail address: imtak-ryk@umin.ac.jp (T. Inoue)

13 November 2013

Available online 25 December 2013